

THE ANTITUMOR ACTIVITY OF ELEMENE IS ASSOCIATED WITH APOPTOSIS

Yang Hua 杨骅 Wang Xianping 王仙平 Yu Linlin 郁琳琳 Zheng Shu 郑树

Cancer Institute, Zhejiang Medical University, Hangzhou 310009

Elemene, isolated from the Chinese medicinal herb *Rhizoma Zedoariae* was shown to exhibit antitumor activity *in vitro* and *in vivo* to human and murine tumor cells. This novel antineoplastic agent has been demonstrated to have substantial clinical activity against various tumors. In this paper, the mechanisms of antitumor activity of elemene are reported. The *in vitro* effect of elemene on the growth of leukemia cells was evaluated by MTT assay. The IC_{50} values of elemene for promyelocytic leukemia HL-60 cells and erythroleukemia K562 cells were found to be 27.5 $\mu\text{g/ml}$ and 81 $\mu\text{g/ml}$, respectively, while IC_{50} for peripheral blood leukocytes (PBL) was 254.3 $\mu\text{g/ml}$. The inhibitory effect of elemene on proliferation of HL-60 cells was associated with cell cycle arrest from S to G_2M phase transition and with induction of apoptosis. The apoptosis of tumor cells was confirmed by DNA ladder formation on gel electrophoresis and characteristic ultrastructural alternations. These results indicate that induction of apoptosis contributes to the mechanisms of antitumor activity of elemene.

Key words: Elemene, Apoptosis, Leukemia

β -elemene (1-methyl-1-vinyl-2, 4-diisopropenyl-cyclohexane) is a naturally occurring compound that can be isolated from the traditional Chinese medicinal herb, *Rhizoma Zedoariae*, native to South China. Elemene has been shown to have variety of pharmacological effects in animal experiments and

clinical trials.¹⁻³ This recent antineoplastic agent has been demonstrated to have substantial clinical activity in treatment of various tumors, but less side-effects and no signs of bone marrow suppression,⁴⁻⁶ the mechanism of which is unclear. In this paper, we determine if the growth inhibitory effects of elemene on human leukemia cells involved in the process of proliferation (cell cycle progression) and/or cell death (apoptosis).

MATERIALS AND METHODS

Chemicals

ELEMENE EMULSION INJECTION (elemene) was originally provided by Dalian Jin Gang Pharmaceutical Co., Ltd. (Dalian, P. R. China). DAPI and other laboratory reagents were purchased from Sigma (St. Louis, MO. USA.). Cisplatin (CDDP) was purchased from Jinzhou Pharmaceutical factory (Jinzhou, P.R. China).

Cell Culture and Treatment

The human promyelocytic HL-60 cells were obtained from Shanghai Institute for Cell Biology, Academia Sinica. The human leukemia K562 cells were kindly provided by Dr. Steinmann (Kiel University, Germany). HL-60 and K562 cells were maintained in RPMI 1640 medium supplemented with 10% fetal bovine serum in our lab. Human peripheral blood mononuclear cells (PBMCs) were isolated from

the healthy donors using Ficoll-Hypaque gradient centrifugation. Cultures of 5×10^4 /ml were generally incubated with elemene, which was diluted in culture medium to obtain the desired concentration. Cells were incubated with various concentration of elemene for 2 to 48 hours. The same aliquot of drug-free emulsion along was added to the control cells.

Cytotoxicity and MTT Assay

IC₅₀ value calculation for each cell line and PBMNs were determined by MTT assay as described.⁷ For MTT assay, cells were seeded at 5×10^4 cells/well and incubated overnight, then treated with equal volumes of medium containing elemene final concentrations from 0.01 to 320 µg/ml. All experiments were repeated at least twice, each in duplicate.

Flow Cytometry Analysis

Cell sample preparation and DAPI staining for flow cytometry analysis were performed according to the method as reported previously.^{8,9} Cell cycle distribution was determined using a CA-II instrument (Partec, Germany). Resulting DNA distribution were then analyzed for proportion of cells in apoptosis. Data was analyzed by Multicycle software (Phoenix Flow System, San Diego, CA).

Electronic Microscope Examination

HL-60 cells treated for 24 hours with different concentrations of elemene. Cell samples were conventionally fixed.¹⁰ Cells were examined and photographed using Philips EM410 transmission electron microscope (TEM).

Agarose Gel Electrophoresis of Apoptotic DNA

Cells were treated with various concentrations of elemene then harvested and washed with phosphate-buffered saline (PBS). 1×10^6 Cell pellets were suspended in 1 ml of lysis solution containing 5 mmol/L Tris-HCl pH 8, 0.25% NP-40, 1 mmol/L EDTA, 10mg/ml RNase at 37 °C for 1 hour. 25 µl of proteinase K (20 mg/ml) was added to the above suspension for 1 hour at 37 °C. The supernatant of the cell lysat was analyzed by electrophoresis in a 1.5% agarose gel containing 0.2% µg/ml ethidium bromide

(EB), and visualized under UV illumination, recorded by a IS-1000 Digital Image System.¹¹ Dexamethasone-induced apoptotic DNA of rat thymus cells was used as positive control in agarose gel electrophoresis.

RESULTS

Inhibitory Effect of Elemene on the Proliferation of Human Leukemia Cells

Cell proliferation of leukemia cells treated with elemene were evaluated by MTT assay. When leukemia cells in culture were treated with elemene for 72 hours, the growth of HL-60 leukemia cells and K562 leukemia cells was significantly inhibited. The IC₅₀ values of elemene for HL-60 and K562 cells were found to be 27.5 µg/ml and 81 µg/ml, respectively, while treated with elemene for 72 hours, IC₅₀ for peripheral blood leukocytes (PBL) from normal blood donor was 254.3 µg/ml.

Effect of Elemene on Cell Cycle Progression of Tumor Cells

When HL-60 cells treated with 10 µg/ml of elemene, the proportion of cells at G₂ M phase was relatively decreased for 4 hours treatment. However, during 24 hours to 48 hours treatment, no significant effect of 10 µg/ml elemene on cell cycle could be observed. When treated with 20 µg/ml elemene, the proportion of cells at G₂ M phase was markedly decreased and dependent on the time of drug exposure. In the histogram, the distinct apoptotic feature of sub-G1 peak (Ap peak) were observed (Figure 1). The percentage of apoptotic cells for the treatment of 4, 24, 48 hours were 41.5%, 35.3% and 47.7%, respectively. When the dose of elemene increased to 40 µg/ml, total cells were undergoing death completely beginning from 4 hours treatment. Cell cycle analysis by flow cytometry revealed that elemene has great influence on cell cycle progression of HL-60 cells and its effect is dependent on the dose and time of drug exposure. The inhibitory effect of elemene on proliferation of HL-60 cells was associated with cell cycle arrest from S to G₂ M phase transition and reduced the proportion of cells at the G₂ M phase, consequently inhibiting mitosis of tumor cells. On the other hand, the elemene-treated tumor cells were rapidly induced undergoing apoptosis.