

Original Article

Health Economic Assessment for Screening of Gastric Cancer in A High Risk Population in Northeastern China

Ling Zhou¹, Peng Guan¹, Li-ping Sun², Qin-cheng He¹, Yuan Yuan², Bao-sen Zhou^{1*}

¹Department of Epidemiology, School of Public Health, ²Cancer Institute, First Affiliated Hospital, China Medical University, Shenyang 110001, China

DOI: 10.1007/s11670-011-0021-7

© Chinese Anti-Cancer Association and Springer-Verlag Berlin Heidelberg 2011

ABSTRACT

Objective: To assess economic cost-effects for the screening programs of gastric cancer in a high risk population in northeastern China.

Methods: The data were collected from November 2001 to December 2003. The multi-stage sampling to define the screening group and the control group was applied in this study. Two stage screening programs were used in the study. An epidemiological survey and serum PG test were carried out in the first stage. The endoscopy and pathological examination were performed in the second stage screening. Effectiveness was assessed by the increased quality adjusted life-year (QALY) because of reduced gastric cancer deaths in screening.

Results: A total of 27,970 participants (n=7,128 screening group, n=20,842 control group) were enrolled in the survey. Twenty nine gastric cancer cases were detected in the screening group with 20 cases in the early stage and 9 cases in the advanced stage, respectively. Eighty six gastric cancer cases were detected in the control group, all of whom were in the advanced stage and had died before the study finished. The screening and treatment of 29 cases cost \$152,227 and \$5,249 per each case, respectively. The costs were \$459 to gain per QALY.

Conclusion: The screening program of gastric cancer used in our study is an economic and society-beneficial measure to detect gastric cancer in high risk area. The methods fit China's present economic development level.

Key words: Gastric cancer; Screening; Economics

INTRODUCTION

Gastric cancer is the fourth most common cancer and the second leading cause of cancer-related death worldwide, especially in Central Europe, South America, and Asia^[1-3]. In China, the incidence of gastric cancer is among the highest of the world, with an estimated 160,000 deaths each year^[4,5]. Gastric cancer detected at a symptomatic stage often has poor survival and high recurrence. Early gastric cancer is typically small and asymptomatic, and the low mortality from gastric cancer is mainly due to early detection. When diagnosed at an early stage, 5-year survival rates for gastric cancer exceed 90%, while for an advanced gastric cancer, 5-year survival rates are below 50%^[6-8]. Therefore, early detection and treatment is an important approach to achieve good prognosis and reduce death from gastric cancer^[9].

Despite the high incidence rate of gastric cancer in China, there is no nationwide screening program to date. Early detection of gastric cancer only relies on the opportunistic screening^[9]. A series of comprehensive

preventive methods have been applied since 1997 in China, including primary and secondary prevention, especially in high-risk populations of gastric cancer.

Zhuanghe County is located in of Liaoning Province with a total population of 900,000, and in the seaside of the Huang Sea. The incidence and mortality of gastric cancer in this county are higher than the average level of China^[10]. From 1996 to 2000, the gastric cancer death counted 8.25% of all causes death, and 38.94% of all cancer death in Zhuanghe County^[11]. The investigations of China Medical University have worked in Zhuanghe for more than 20 years and established a comprehensive gastric cancer prevention network^[12].

Serum pepsinogen (PG) has been found to be a marker of gastric mucosal status, and associated with gastric cancer^[13]. Measurement of serum PG is a popular non-invasive serological screening test for identifying individuals with extensive gastric atrophy that warrants more intensive surveillance for high-risk of gastric cancer^[14]. Endoscopy and pathological examinations have been increasingly used for gastric cancer diagnosis as a golden standard because of a high detection rate.

The two-stage screening program has good sensitivity and specificity in the detection of gastric cancer. Therefore, the relative costs, benefits and long term outcome are important factors to consider when evaluating the public

Received 2010-07-29; Accepted 2010-11-26

This study was supported by National Key Technologies R&D Program of China during the 10th Five-year Plan Period (No. 2004BA703B04-02).

*Corresponding author.

E-mail: bszhou@mail.cmu.edu.cn

health relevance of this intervention. This study aims to evaluate the economic cost-effects of the screening program for gastric cancer in a high-risk population in the Northeast China.

MATERIALS AND METHODS

Definition of Screening Population

The multi-stage sampling to define the screening group and the control group was applied in this study. Firstly, villages were stratified into two parts based on the landform, seaside or hill. Secondly, then each part was divided into two groups, the screening group and the control group, using the method of complete random cluster sampling.

Among the 28 sampled villages, 15 villages were assigned to the screening group and 13 villages were assigned to the control group. There were no difference between two groups in the geographical environment, the humanity environment and other basic characteristics.

Screening

The design of the screening is presented in Figure 1.

The target population was those who had the family history of gastric cancer, and /or were over 35 years old and had the history of gastric illness, or had evident gastric illness symptoms. A two-stage screening program was conducted. An epidemiological survey and serum PG test were carried out in the first stage. If the serum PG I/II ratio was less than 9 and the participant was younger than 50 years old, or the ratio was less than 7.5 and subject was over 50 years old, or PG I was less than 70 µg/L in male, or PG I was less than 60 µg/L in female, the participant would receive the second stage screening with endoscopy and pathological examination.




Figure 1. The screening trial sketch

Calculation of Costs

The direct costs for PG tests, endoscopy/pathology and labor in the screening were estimated by the program. The

data for the other expenditure for gastric cancer treatment were obtained from hospital finance department, patients and relatives. The treatment costs included out-patient cost, drug cost, surgery cost and hospitalization cost. Because the screening program had certain time span and the medical cost had largely fluctuated, costs and benefits were discounted to the price level in 2001. The discount rate was defined by national retail price general index and resident consumption price general index. A 5% discount rate was adopted in our study.

Effectiveness can be seen from the increased quality adjusted life-year (QALY) because of reduced gastric cancer deaths in screening. We used the World Health Organization Quality of Life (WHOQOL)- BREF to measure the survival quality of the gastric cancer patients, and then to estimate the effectiveness.

Data Collection and Analysis

The data were collected from November 2001 to December 2003. The first stage screening took place in the appointed clinic in each village, while the second stage screening was carried out in the Zhuanghe Central Hospital and Zhuanghe Hospital of Traditional Chinese Medicine. The health workers who provided screening had been trained. Cost data were obtained from hospital finance departments, patients and their relatives. Annual income was acquired from the local government finance department of screening villages. Quantitative data were input and analyzed using SPSS 16.0.

Ethical Issues

This study was approved by the Ethics Committee of China Medical University and implemented under the full knowledge and consent by participants, who were assured of the security of the information. Written, informed consent was received from all study participants before the screening.

RESULTS

A total of 27 970 participants (n=7 128 screening group, n= 20 842 control group) were included in the survey. Of them, 50.86% were males while 49.14% were females in the screening group, and 50.54% were males while 49.46% were females in the control group (Table 1).

Table 1. Composition of the samples in high risk area of Zhuanghe, China, November 2001 to December 2003

Screening group	Male	Female	Total
Screening group	3 625	3 503	7 128
Control	10 533	10 309	20 842
Total	14 158	13 812	27 970

Screening Result

Twenty nine gastric cancer cases were detected in the screening group. Twenty cases were in the early stage, and 9 cases were in the advanced stage. Eighty six gastric cancer cases were founded in the control group, who were all in the advanced stage and had died before the study finished.